



ગ્રામીણ કૃષિ મૌસમ સેવા

કૃષિ હવામાન સેવા

નવસારી કૃષિ યુનિવર્સિટી

નવસારી - ૩૮૬૪૫૦, ગુજરાત

E-મેઈલ:- aasnaysari@gmail.com



(નવસારી કૃષિ યુનિવર્સિટી અને ભારત મૌસમ વિજ્ઞાન વિભાગ દ્વારા પ્રસારિત)

તા. ૨૪/૦૮/૨૦૨૧

કૃષિ હવામાન બુલેટીન નં : ૫૧/૨૦૨૧ - ૨૦૨૨

-- નવસારી જિલ્લો --

પાછળનાં સમયનું હવામાન (તા. ૨૧ સપ્ટેમ્બરથી ૨૪ સપ્ટેમ્બર - ૨૦૨૧)

અનું. નં.	હવામાન પરિબળો	Value
૧.	વરસાદની શ્રેણી (મી.મી.)	૮.૦-૯૧.૦
૨.	મહત્તમ તાપમાનની શ્રેણી (સે.ગ્રે.)	૨૭.૮-૩૧.૫
૩.	લધુત્તમ તાપમાનની શ્રેણી (સે.ગ્રે.)	૨૨.૨-૨૩.૮
૪.	ભેજની શ્રેણી (%)	૮૫-૯૦૦
૫.	પવનની ગતિની શ્રેણી (કિ.મી./કલાક)	૨.૦-૩.૫
૬.	વાદળની સ્થિતિની શ્રેણી (ઓક્ટા)	૭-૮
૭.	સુર્યપ્રકાશિત સમયની શ્રેણી (કલાક)	૦.૦-૪.૩
૮.	પાછળના પાંચ દિવસમાં સંચિત વરસાદ (મી.મી.)	૮૪.૦
કુલ વરસાદ		૧૫૧૭.૦

આઈ. એમ. ડી. દ્વારા પ્રાપ્ત હવામાન આગાહી (તા. ૨૫ સપ્ટેમ્બરથી ૨૮ સપ્ટેમ્બર - ૨૦૨૧)

અનું. નં.	હવામાન પરિબળો	દિવસ-૧ (૨૫/૦૮)	દિવસ-૨ (૨૬/૦૮)	દિવસ-૩ (૨૭/૦૮)	દિવસ-૪ (૨૮/૦૮)	દિવસ-૫ (૨૯/૦૮)
૧.	વરસાદ (મી.મી.)	૫૫	૬૦	૫૦	૬૫	૫૫
૨.	મહત્તમ તાપમાન (સે.ગ્રે.)	૩૧	૩૧	૩૦	૩૧	૩૧
૩.	લધુત્તમ તાપમાન (સે.ગ્રે.)	૨૬	૨૫	૨૫	૨૬	૨૬
૪.	મહત્તમ ભેજ (%)	૮૨	૮૨	૮૩	૮૩	૮૨
૫.	લધુત્તમ ભેજ (%)	૭૬	૮૩	૮૬	૮૨	૮૫
૬.	પવનની ગતિ (કિ.મી./કલાક)	૧૩	૧૬	૧૫	૧૫	૧૨
૭.	પવનની દિશા (ક્રમપાસ પોર્ટન્ટ)	૨૦૭	૨૪૩	૨૨૫	૧૮૦	૧૮૦
૮.	વાદળની સ્થિતિની (ઓક્ટા)	૬	૭	૭	૮	૮

હવામાન સારાંશ / ચેતવણી

- આઈ. એમ. ડી., દ્વારા મળેલી હવામાનની આગાહી મુજબ આગામી પાંચ દિવસમાં આકાશમાં વાદળાયું રહેશે અને છુટા છવાયા વિસ્તારમાં મધ્યમથી ભારે વરસાદ પડવાની સંભાવના છે.
- મહત્તમ તાપમાન ૩૦ થી ૩૧ સે. જ્યારે લધુત્તમ તાપમાન ૨૬ થી ૨૮ સે. આસપાસ રહેવાની સંભાવના છે.
- હવામાં ભેજનું પ્રમાણ ૭૬ થી ૮૨ ટકા વચ્ચે રહેવાની સંભાવના છે.
- પવનની દિશા દક્ષિણ-પ્રિયમ રહેશે જેની સરેરાશ ગતિ ૧૨ થી ૧૬ કિ.મી./કલાક રહેવાની શક્યતા છે.
- તારીખ ૦૫ ઓક્ટોબર, ૨૦૨૧ સુધી, લાંબા ગાળાની આગાહી મુજબ સામાન્ય થી ઉપર (૧૧.૨૬ મિમી.) વરસાદ પડવાની સંભાવના છે.

-:: દક્ષિણ ગુજરાત ભારે વરસાદ વિસ્તાર ::-

<p>પાક</p> <p>ચોમાસુ ડાંગર</p> <p>કૂલ / દાણા ભરાવા અવસ્થા:</p> <ul style="list-style-type: none"> • કૂલ અવસ્થા અને દાણા ભરાવાની અવસ્થા દરમ્યાન ૪-૫ સે. મી. જેટલું પાણી ભરી રાખવું કાપણી : • કાપણીના ૧૦-૧૫ ટિવસ ફેલા ક્યારીમાંથી પાણી નિતારી દેવું • વરસાદ ની સંભાવના ન હોય ત્યારે કાપણી કરવી. • કાપણી કર્યો બાદ સૂર્ય તાપમાં સૂક્કવી દાણા છૂટા પાડવા અને દાણાનો ભેજ ઓછો થાય તાં સુધી તપાવી સંગ્રહ કરવો. <p>શેરડી</p> <p>વાનસ્પતિકવૃદ્ધિ અવસ્થા:</p> <ul style="list-style-type: none"> • શેરડીના વધેકો માટે કાર્બોફ્યુરાન ઉજી ઉજી કી.ગ્રા./લે. જમીનમાં આપવું હવામાન ચોખ્ખુ હોય ત્યારે છંટકાવ કરવો... • શેરડીની સફેદમાખીના નિયંત્રણ માટે બેતરમાં ભરાયેલ પાણીનો નિકાલ કરવો. • શેરડીની નીચેની સૂક્કી પાતરી ફ, ઉ અને ઈ માસે બેરવી નાંખવી. 	
---	--

અન્ની નાવસરી

	<p>નવી રોપણી માટે જમીન તૈયારી:</p> <ul style="list-style-type: none"> શેરડીની નવી રોપણી માટે યોગ્ય બેજની પરિસ્થિતિમાં જમીન એયર કરવી. શેરડીની રોપણી ઓક્ટોબર-ફેબ્રુઆરી માસ દરમિયાન કરવી. ખેડ વધતે ખાતર/કમ્પોસ્ટ જમીનમાં ભેગવું. રોપણી પહેલા પાયાનું ખાતર આપવું. રોપણી માટે કુમળૂ અને તંડુરસ્ત તમારી પસંદની જાતનું બિયારણ વાપરવું. રોપણી માટે બે લાઈન વર્ચ્યે ૮૦ થી ૧૨૦ સેમી. અંતર રાખી બે આંખના કટકાને બીજમાવજત આપી એક પછી એક કટકા ચાસમાં રહે તે રીતે રોપણી કરવી. એક આંખના રોપનું વાવેતર કરવાનું હોય તો લાઈનમાં બે રોપા વર્ચ્યે ૪૫ થી ૬૦ સેમી. અંતર રાખી રોપણી કરવી. રોપણી બાદ પિયત આપી ઉ દિવસમાં એટ્રોપ્લીન ૫૦ ડબલ્યુ પી પ્રી ઇમરજન્સ નીંદુનાશક નો છંટકાવ હવામાન ચોખ્ખુ હોય ત્યારે કરવો.
ભૂંડા	<ul style="list-style-type: none"> ખેતરમાં પાણીની ભરાવો ટાળવો.
શક્કભાજી	<ul style="list-style-type: none"> નોવેલ ૧ લિટર/૧ વીધામાં આપવું. નોવેલ પ્રવાહી ખાતર ૧૫૦ મી.લી./૧૫ લિટર પાણીમાં નાખી ઊભા પાકમાં છંટકાવથી પાક ઉત્પાદન વધારી ગુણવત્તાસભર પાક મેળવી શકાય છે અને હવામાન ચોખ્ખુ હોય ત્યારે છંટકાવ કરવો. નોવેલ કાર્બનિક પ્રવાહી પોષક ભાવ રૂ. ૧૨૦/- પ્રતિ લિટર છે. નોવેલ એસ તથા નોવેલ પ્રાઇમ નો ભાવ રૂ. ૨૦૦/- પ્રતિ લિટર છે.
રોગણુ	<ul style="list-style-type: none"> ધરુ ૨૮-૩૦ દિવસનું થાય ત્યારે ફેરરોપણી કરવી.
ટામેટાં	<ul style="list-style-type: none"> ધરુ ૨૫-૩૦ દિવસનું થાય ત્યારે ફેરરોપણી કરવી.
મરચાં	<ul style="list-style-type: none"> મરચીનું ધરુ લગભગ ઉપ થી ૪૦ દિવસે રોપવા લાયક બને છે.
ચોળી	<ul style="list-style-type: none"> ચોમાસાની ઋતુ માટે વાવણી કરી શકાય છે. ચોળીનું વાવેતર બે હાર વર્ચ્યે ૩૦ × ૪૫ સે.મી. અંતર રાખી કરવું. એક લેક્ટરના વિસ્તાર માટે ૧૨-૧૫ કિ.ગ્રા. બિયારણની જરૂર પડે છે.
વેલાવાળા શક્કભાજી પાકો	<ul style="list-style-type: none"> વાવણી બાદ ૧૦ થી ૧૨ દિવસે કરબળીથી આંતર ખેડ કરવી. ન્યારબાદ ૨ થી ૩ વખત હાથથી નીંદામણ કરતાં રહેવું.
સુરણુ	<ul style="list-style-type: none"> ખેતરમાં પાણીની ભરાવો ટાળવો.
રતાળુ	<ul style="list-style-type: none"> ખેતરમાં પાણીની ભરાવો ટાળવો.
તાનિયા	<ul style="list-style-type: none"> ખેતરમાં પાણીની ભરાવો ટાળવો. “પાતરા” નામની વાનગી બનાવવા માટે કુમળા પાનની વીણી કરતા રેહવું જોઈએ.
આંખા	<p>સામાન્ય સલાહ:</p> <ul style="list-style-type: none"> આંખાના મેઢની ઈયણો થડમાંથી શોધીને મારી નાખો અને કાણાંની અંદર ક્લોરેપાયરોફોસ ૨૫ મી.લી./લી. ટ્રાવણ નાંખી કાદવથી કાણાં બંધ કરી દેવા હવામાન ચોખ્ખુ હોય ત્યારે છંટકાવ કરવો.

	<ul style="list-style-type: none"> આંબા નવી વાડી બનાવવા તંદુરસ્ત અને રોગમુક્ત કલમો અધિકૃત અને સરકારી નર્સરી માંથી મેળવવી જોઈએ. નવી કલમો માટે તૈયાર કરેલ ખાડામાં ૧૦ કિગ્રા. છાણિયું ખાતર, ૧ કિગ્રા. સુપર અને ૧ કિગ્રા. પોટાશ ખાતર આપવું ઉધૃઠ અને કીડીનો ઉપદ્રવ કલમોમાં જોવા મળે તો ૫૦ મિલી પ્રતિ ૧૦ લીટર લોરપાયરેફોસ ની માવજત ખાડા પૂરતી વખતે કરી દેવી હવામાન ચોખ્ખુ હોય તારે છંટકાવ કરવો. આંબાવાડીમાં પુષ્ટ જાડ દિંદ ૧૦૦ કિગ્રા. છાણિયું ખાતર, ૧૦૦ ગ્રામ યુરિયા અથવા ૧.૮ કિગ્રા. સુપર અને ૧.૨ કિગ્રા. મુરેટ ઓફ પોટાશ ખાતર ચોમાસામાં આપવું હવામાન ચોખ્ખુ હોય તારે છંટકાવ કરવો. આંબાવાડીયામાં રાસાયણિક ખાતર આપ્યાના ૧૫ દિવસ પહેલા અથવા પછી બાયોફાઈલ્ડર જેવા કે એઝોટોબેક્ટર, ૫૦ મિલી., ફોસ્ફેટ સોલ્યુબિલિઝિંગ બેક્ટેરિયા ૫૦ મિલી., પોટાશ મોબિ લિઝિંગ બેક્ટેરિયા ૫૦ મિલી. અને ટ્રાયકોર્ટમાં ૧૦૦ ગ્રામ. પ્રતિ જાડ આપવું હવામાન ચોખ્ખુ હોય તારે છંટકાવ કરવો. ડાળીનો સુકારો : રોઝીએ ડાળીઓ કાપી, કપાયેલા ભાગ પર ૧૦ ટકા બોર્ડકસ પેસ્ટ લગાડવું તેમજ કાર્બન્ડાલ્જિમ ૫૦ ઉભલ્યુ પી ૧૦ ગ્રા. પ્રતિ ૧૦ લીટર પાણીમાં મિશ્ર કરી છંટકાવ કરવો હવામાન ચોખ્ખુ હોય તારે છંટકાવ કરવો. આંબાવાડીમાં ભરાયેલ પાણીનો નિકાલ કરવો.
ચીકુ	<p>સામાન્ય સલાહ:</p> <ul style="list-style-type: none"> બીજા હન્તામાં ભલામણ કરેલ ખાતરનો ૫૦ ટકા ભાગ પ્રમાણે રાસાયણિક ખાતર એમોનિયમ સફ્ટેક્ટ ૨.૫ કિગ્રા. અથવા યુરિયા ૧.૩ કિગ્રા., સિંગલ સુપર ફોસ્ફેટ ૧-૨ કિગ્રા., મુરેટ ઓફ પોટાશ ૦.૫ કિગ્રા., છાણિયું ખાતર ૫૦ કિગ્રા. અને બાયોફાઈલ્ડર (૫૦ મી.લી. - ૫૦ મી.લી. એઝોટોબેક્ટર, ફોસ્ફેટ સોલ્યુબિલિઝિંગ બેક્ટેરિયા અને પોટાશ મોબિ લિઝિંગ બેક્ટેરિયા) આ મહિનામાં આપવું હવામાન ચોખ્ખુ હોય તારે છંટકાવ કરવો.
પશુપાલન	<p>સામાન્ય સલાહ:</p> <ul style="list-style-type: none"> પશુઓને દૂધ ઉત્પાદન અને નિભાવની જરૂરીયાત ધ્યાનમાં લઈ પૂરતા પ્રમાણમા સમતોલ આહાર આપવો. તેમજ પશુઓના રહેઠાળ ચોખ્ખા રાખવા, દિવસ દરમયાન સમયાંતરે મળ-મૂતર નો નિકાલ કરવો. પશુઓને યોગ્ય આહારમા જરૂરીયાત પ્રમાણે ખનિજયુક્ત મિશ્રણના સમાવેશ કરવો જોઈએ. ચાદ્ર કટરથી લીલા તથા સૂકી ઘાસચારાને ટુકડા કરીને જ નીરણ કરવું. દુધાળા પશુઓના આઉને દૂધ દોહન પહેલા અને પછી ઝીક ઓક્સાઈડ અથવા બોરીક પાઉડર અથવા પોટેશીયમ પરમેગેનેટ ના ટ્રાવણુથી ભરાબર સાફ્ટ કરવા. પશુઓનાં ઈતરરીના નિયંત્રણ માટે ડેવટામેથીન અથવા એમીટાઇર ૨ મી.લી./લિટર પાણીમાં નાખને છાંટવી (દવાની બોટલ ઉપર લખેલી સૂચનાનો અમલ કરવો).

	<ul style="list-style-type: none"> પશુઓ જ્યારે દીચ્છે ત્યારે પાણી પી શકે તેવી વ્યવસ્થા ઉભી કરવી જોઈએ અથવા ઓછામાં ઓછું દિવસમાં ચારથી પાંચ વખત જરૂરીયાત મુજબ પાણી પીવડાવવું જોઈએ. ચોમાસાની શરૂઆત થની હોઈ પશુઓના રહેઠાળમાં વરસાઈ પાણીનો ભરાવો ના થાય તે પ્રમાણે કોઢ/ગમાણના ભોયાતળિયાનો ઢોળાવ અપાવો. કોઢ/ગમાણની ઉચ્ચાઈ ૧૦ ફૂટથી વધારે રાખવી, જેથી હવાની હેરફેર સરળતાથી થઈ શકે, બફારો/ગરમી ઓછી થાય અને ભોયાતળિયાનો ભાગ સુકો રહે. સતત પુષ્કળ વરસાઈમાં નાના વાષ્પરડા-પાઢીયા કે અન્ય પુષ્ણત પશુ સતત લાંબા સમય સુધી ભીજ્ય નહિ તે માટે તાડપત્રી/પનરાનો શેડ કે પાકી ગમાણ હેઠળ રાખવા. ખુબ વીજળી અને વરસાઈના સમયે પશુ વૃક્ષ નીચે બાંધવા નહિ.
વનીકરણ	<p>કૃષિ વનીકરણ : વાવેતર :</p> <ul style="list-style-type: none"> પ્રથમ બને ભીજ વર્ષના વાવેતર દરમ્યાનમાં નિંદણ અને સફાય કરવી જોઈએ. નવા ટ્રાન્સપ્લાન્ટ થયેલા રોપાઓ ખાસ કરીને જળસંચય અને છોડના રક્ષણથી સંભાળી શકાય છે.
ફીશરીજ	<p>સામાન્ય સલાહ :</p> <ul style="list-style-type: none"> મીઠા પાણીના મ ત્સય મીઠા પાણીના જુંગા ઉત્યાદન કરતા ખેડૂતોને જણાવવાનું કે , આ સમયગાળા દરમ્યાન પૂરતી માત્રામાં ખોરાક આપવોકારણ કે આ વાતાવરણ માછલી/મીઠા પાણીના જુંગા માટે અનુકૂળ છે. તળાવમાં પાણીનું લેવલ જાળવવું અને વધુ વરસાઈ આવે ત્યારે તળાવના આઉટલેટ થોડી માત્રામાં ખોલવા તેથી પાણીનું લેવલ જણાવાયેલું રહે. તળાવના તળિયાનું અને સપાટીનું નાપમાન સરખર કરવા એરેટર અથવા પંખ ચલાવવા.

NAVSARI AGRICULTURAL UNIVERSITY

❖ એન્ગ્રોમેટ સલાહકાર સેવા બુલેટીન ફિઝિટીના સભ્યો :

- ડૉ. એચ. એમ. વિરદીયા, સંશોધન વૈજ્ઞાનિક (સશ્ય વિજ્ઞાન)
- ડૉ. એલ. વી. ઘેરીયા, સહ પ્રાધ્યાપક (વનસ્પતિ કીટકશાસ્ત્ર)
- ડૉ. કે. બી. રાખોલીયા, સહ પ્રાધ્યાપક (વનસ્પતિ રોગશાસ્ત્ર)
- ડૉ. એસ. એન. સરવૈયા, પ્રધ્યાપક (શક્કભાજ વિજ્ઞાન)
- ડૉ. વાય. એન. ટેલે, સહ પ્રાધ્યાપક (કૃળ વિજ્ઞાન)
- ડૉ. વી. આર. પટેલ, મદદનીશ પ્રાધ્યાપક (પશુપાલન)
- ડૉ. દિવેશ્વર નાયક, મદદનીશ પ્રાધ્યાપક (વનીકરણ)
- ડૉ. એમ. આર. પટેલ, મદદનીશ પ્રાધ્યાપક (ફીશરીજ સાયન્સ)
- ડૉ. વિભા ટાક, રીસર્ચ એસોસીયેટ (જ.કે.એમ.એસ.)
- ડૉ. પી. કે. પરમાર , મદદનીશ પ્રાધ્યાપક (કૃષિ હવામાનશાસ્ત્ર) અને પ્રિન્સપાલ નોડલ ઓફિસર (જ.કે.એમ.એસ.)



Gramin Krishi Mausam Sewa

Agricultural Meteorological Cell
Navsari Agricultural University
Navsari – 396450, Gujarat
E-mail :- aasnnavsari@gmail.com



(Jointly issued by Navsari Agricultural University & India Meteorological Department)

DATE OF ISSUE: 24/09/2021

Agromet Advisory Bulletin No: 51/2021 - 2022

-- Navsari District --

Weather summary of preceding week 21 September to 24 September - 2021

Sr. No.	Name of Weather Parameters	Value
1.	Range of Rainfall (mm)	8.0-61.0
2.	Range of Maximum Temperature (°C)	27.8-31.5
3.	Range of Minimum Temperature (°C)	22.2-23.8
4.	Range of Relative Humidity (%)	85-100
5.	Range of Wind Speed (km/hr)	2.0-3.5
6.	Range of Cloud Cover (Okta)	7-8
7.	Range of bright sunshine hours (hrs)	0.0-4.3
8.	Last five days cumulative rainfall (mm)	94.0
Total Rainfall (mm)		1517.0

IMD based Weather forecast for 25 September to 29 September -2021


Sr. No.	Weathers Parameters	Day - 1 (25/09)	Day - 2 (26/09)	Day - 3 (27/09)	Day - 4 (28/09)	Day - 5 (29/09)
1.	Rainfall (mm)	65	90	60	95	65
2.	Maximum temperature (°C)	31	31	30	31	31
3.	Minimum Temperature (°C)	26	25	25	26	26
4.	Max. Relative Humidity (%)	92	92	93	93	92
5.	Min. Relative Humidity (%)	76	83	86	82	85
6.	Wind Speed (km/hr.)	13	16	15	15	12
7.	Wind Direction (Compass point)	207	243	225	180	180
8.	Total Cloud Cover (Okta)	6	7	7	8	8

Weather Summary/ Alert	<ul style="list-style-type: none"> As per IMD weather forecasts over the next five days, the sky will be cloudy and there is chance of moderate to heavy rainfall occurs at isolated places. The Maximum temperature 30 to 31 °C. when the Minimum temperature is 25 to 26 °C. likely to be around. The humidity in the air is likely to be between 76 and 93 percent. The wind direction will be South-Westerly with an average speed of 12 to 16 km/h. As per Extended range forecast, possibility of rainfall in Gujarat region is likely to be above normal (11.26 mm) up to 05th October, 2021.
-----------------------------------	--

-:: South Gujarat Heavy Rainfall Zone ::-

General Advisory <ul style="list-style-type: none"> • Due to pandemic of CORONA (COVID-19), Farmers should maintain social distancing and safety precaution while handling farm machines and labour in the field. • Farmers are advised to postpone the activity of spraying chemical and applying fertilizer • Farmers are advised to postpone the different activities of farm operations. • Farmers are advised to arrange drainage facility to remove excess rain water from field. • Fisher man is advice not to venture into sea area during this heavy rainfall time. • Download the MEGHDOOT mobile application, We will provide forecast to farmers relating to temperature, rainfall, humidity, wind speed and direction which play critical roles in agricultural operations and how to take care of the crops and livestock. Click on the following link to download the MEGHDOOT application. Android users: https://play.google.com/store/apps/details?id=com.aas.meghdoot IOS users: https://apps.apple.com/in/app/meghdoot/id1474048155
--

NOAA/VIIRS/BLENDED NDVI Composite ending on Week no 37
(10.09.2021 to 16.09.2021) over Agricultural region



Agricultural vigour is good over isolated pockets of the state.

Crop	Weather based Agro-Advisory
Kharif Paddy	<p>Flowering stage/Grain filling:</p> <ul style="list-style-type: none"> • Keeps 4-5 cm water in field during flowering and grain filling stage. <p>Harvesting:</p> <ul style="list-style-type: none"> • Drain the water from the kyari 10-15 days prior to the harvesting. • Harvest the crop when there is no possibility of rainfall and sky is clear. • After harvesting properly dry the crop and threshed the grains, also dry the grains to remove moisture before storage.
Sugarcane	<p>Grand growth stage/Maturity:</p> <ul style="list-style-type: none"> • Application of <i>Carbofuran 3G</i> @ 33 kg/ha. in soil for control of sugarcane borer in clear weather condition. • Drain stagnant water from the field for management of sugarcane whitefly. • Detach below dry leaves at 6, 7 & 8 month of planting. <p>Land preparation for new planting:</p>

	<ul style="list-style-type: none"> • New plantation of sugarcane crop should be done after proper preparation of land at proper moisture condition. Deep plough and level the soil. • Planting should be done in October -February. • Apply FYM/ Compost at the time of ploughing. Select healthy, diseases-insect free and immature seed materials of your selected variety after seed treatment. • Planting should be done at the spacing of 90 to 120 cm. 2 budded setts should be used with end to end method of planting. If settling use intra row spacing should be 45 to 60 cm. • Within 3 days after planting and irrigation, spraying of <i>Atrazine</i> 50 WP 2 kg/ha. pre-emergence weedicide should be applied to control weeds in clear weather condition.
Okra	<ul style="list-style-type: none"> • Avoid water logging in the field.
Vegetable Crop	<ul style="list-style-type: none"> • Give Novel 1 lit./Vigha in soil in clear weather condition. • Spraying of Novel 150 ml./15 lit. water on standing crop can increase the crop yield and helps to get quality crop during clear weather condition. • The rate of NOLN (Novel Organic Liquid Nutrient) for 1 lit. is 120 Rs. • NOLN (Novel plus) and NOLN (Novel prime) 200 Rs/lit.
Brinjal	<ul style="list-style-type: none"> • Transplanting should be carried out when the seedlings are 28-30 days old.
Tomato	<ul style="list-style-type: none"> • Transplanting should be carried out when the seedlings are 25-30 days old.
Chilli	<ul style="list-style-type: none"> • Seedlings of chilli become suitable for transplanting in about 35 to 40 days.
Cowpea	<ul style="list-style-type: none"> • Sowing of seed is carried out. • Spacing: 30 x 45 cm. • 12-15 kg seeds are required for one hectare area.
Gourds	<ul style="list-style-type: none"> • Ten to twelve days after sowing, intercultural operations can be carried out with hoe (gardening tool). Then manual weeding can be carried out 2-3 times.
Elephant foot yam	<ul style="list-style-type: none"> • Avoid water logging in the field.
Greater yam	<ul style="list-style-type: none"> • Avoid water logging in the field.
Tannia	<ul style="list-style-type: none"> • Avoid water logging in the field. • Harvesting should be continue for the tender leaves of Tannia for making “paatra”.
Mango	<p>General Advise :</p> <ul style="list-style-type: none"> • Find grubs of stem borer from gallery and holes on the stem and kill them. Do pouring of <i>Chlorpyrifos</i> 20 EC @ 25 ml/ 10 lit. in hole and plug it with mud in clear weather condition. • Procure healthy and disease free planting material from authorized and

	<p>government nursery for the new plantation.</p> <ul style="list-style-type: none"> • Prepare pit for plantation of new grafts by filling 10 kg <i>FYM</i>, 1 kg <i>SSP</i> and 1 kg <i>MOP</i>. Give 50 ml. /10 lit. <i>Chlorpyrifos</i> to manage termite and ants in clear weather condition. • Apply recommended dose of fertilizer per tree @ 700 g <i>Urea</i> or 1.8 kg <i>Ammonium Sulphate</i>, 1.0 kg <i>SSP</i> and 1.2 kg <i>MOP</i> along with 100 kg <i>FYM</i> to adult tree in clear weather condition. • In addition RDF, give biofertilizers viz., <i>Azotobacter</i> 50 ml, <i>PSB</i> 50 ml, <i>KMB</i> 50 ml and <i>Trichoderma</i> 100 g per adult tree in monsoon. Before or after 15 days of fertilizer application in clear weather condition. • Die back disease: Pruning of infected branches, apply 10 % Bordeaux paste and spraying <i>Carbendazim</i> 50 wp, 10gm/10 lit water in clear weather condition. • Improved drainage, remove rain water from new plantation of mango.
Sapota	<p>General Advise :</p> <ul style="list-style-type: none"> • Apply second split of 50% of RDF (<i>Ammonium Sulphate</i> -2.5 kg or <i>Urea</i> 1.3 kg, <i>SSP</i> 1-2 kg and <i>MOP</i> 0.5 kg + <i>FYM</i> 50kg + <i>Azotobacter</i>, <i>PSB</i> and <i>KMB</i> @ 50 ml of each tree) in clear weather condition.
Livestock	<p>General Advise :</p> <ul style="list-style-type: none"> • Provide adequate quantity of balanced ration by considering the milk production and maintenance. • Good quality dry roughage with 15-20 kg green fodder to the animals only after chaffing. • Feed silage & hay materials if available. • Before and after milking udder must be properly clean with zinc oxide or boric powder or potassium permanganate solutions. • For the control of tock and other parasites, spray <i>Deltemethrin</i> or <i>Aemitaz</i> 2 ml in 1 liter of water. • In rainy season, give sufficient slop on the floor of animal shed to avoid water logging. • Height of animal shed should keep minimum 10 feet for aeration to maintain the floor dry and to control humidity. • Don't keep the animal under tree when there is heavy lightning during rain.
Forestry	<p>Agro Forestry : Plantations :</p> <ul style="list-style-type: none"> • Weeding and cleaning should be done during first and second year plantations. • Newly transplanted seedlings may be taken care especially from water logging and plant protection
Fisheries	<p>General Advise :</p>

- Fresh water farmers give proper feed in fish/prawn farming for better growth rate because environment condition is very suitable.
- Increase water depth and maintain water level through pond outlet.
- Mixing the top and bottom water by using the aerators or pumps.

❖ **Members of Agro-Advisory Committee :**

- Dr. H. M. Viradia, Research Scientist (Agronomy)
- Dr. L. V. Ghetiya, Associate Professor (Entomology)
- Dr. K. B. Rakholiya, Associate Professor (Plant Pathology):
- Dr. S. N. Saravaiya, Professor (Vegetable Science)
- Dr. Y. N. Tandel, Associate Professor (Fruit Science)
- Dr. V. R. Patel, Assistant Professor (Animal Husbandry)
- Dr. Dileswar Nayak, Assistant Professor (Forestry)
- Dr. M. R. Patel, Assistant Professor (Fisheries Science)
- Dr. Vibha Tak, Research Associate(GKMS)
- Dr. P. K. Parmar, Assistant Professor (Agricultural Meteorology and Principal Nodal Officer (GKMS))



NAVSARI AGRICULTURAL UNIVERSITY
AMFU_Navsari

જવામાન આગાહી

વિગત	તારીખ	ડાંગ	નવસારી	વલસાડ
વરસાદ (મી.મી.)	૨૫/૦૮/૨૦૨૧	૩૦	૬૫	૬૫
	૨૬/૦૮/૨૦૨૧	૬૫	૬૦	૧૦૦
	૨૭/૦૮/૨૦૨૧	૬૫	૬૦	૬૫
	૨૮/૦૮/૨૦૨૧	૨૦	૬૫	૬૦
	૨૯/૦૮/૨૦૨૧	૪૦	૬૫	૫૦
મહત્તમ તાપમાન (સે.ગ્રે.)	૨૫/૦૮/૨૦૨૧	૩૧	૩૧	૩૧
	૨૬/૦૮/૨૦૨૧	૩૧	૩૧	૩૧
	૨૭/૦૮/૨૦૨૧	૩૦	૩૦	૩૦
	૨૮/૦૮/૨૦૨૧	૩૧	૩૧	૩૧
	૨૯/૦૮/૨૦૨૧	૩૧	૩૧	૩૧
વધુતમ તાપમાન (સે.ગ્રે.)	૨૫/૦૮/૨૦૨૧	૨૬	૨૬	૨૬
	૨૬/૦૮/૨૦૨૧	૨૫	૨૫	૨૫
	૨૭/૦૮/૨૦૨૧	૨૫	૨૫	૨૫
	૨૮/૦૮/૨૦૨૧	૨૬	૨૬	૨૬
	૨૯/૦૮/૨૦૨૧	૨૬	૨૬	૨૬
મહત્તમ ભેજ (%)	૨૫/૦૮/૨૦૨૧	૬૭	૬૨	૬૪
	૨૬/૦૮/૨૦૨૧	૬૭	૬૨	૬૪
	૨૭/૦૮/૨૦૨૧	૬૬	૬૩	૬૪
	૨૮/૦૮/૨૦૨૧	૬૫	૬૩	૬૪
	૨૯/૦૮/૨૦૨૧	૬૬	૬૨	૬૩
વધુતમ ભેજ (%)	૨૫/૦૮/૨૦૨૧	૭૧	૭૬	૭૫
	૨૬/૦૮/૨૦૨૧	૮૦	૮૩	૮૦
	૨૭/૦૮/૨૦૨૧	૮૫	૮૬	૮૪
	૨૮/૦૮/૨૦૨૧	૭૪	૮૨	૮૧
	૨૯/૦૮/૨૦૨૧	૭૩	૮૫	૮૪
પવનની ગતિ(ક્રિ.મી./કલાક)	૨૫/૦૮/૨૦૨૧	૧૦.૫	૧૩.૩	૧૦.૬
	૨૬/૦૮/૨૦૨૧	૧૨.૫	૧૬	૧૩.૩
	૨૭/૦૮/૨૦૨૧	૧૧.૩	૧૪.૭	૧૨.૬
	૨૮/૦૮/૨૦૨૧	૮.૪	૧૫.૧	૧૩.૩
	૨૯/૦૮/૨૦૨૧	૬.૭	૧૨.૨	૧૧.૨
પવનની દિશા (ક્રમપાસ પોઈન્ટ)	૨૫/૦૮/૨૦૨૧	SW	SW	SW
	૨૬/૦૮/૨૦૨૧	SW	SW	SW
	૨૭/૦૮/૨૦૨૧	SW	SW	S
	૨૮/૦૮/૨૦૨૧	SW	S	S
	૨૯/૦૮/૨૦૨૧	SW	S	S
વાદળની સ્થિતિની (ઓક્ટા)	૨૫/૦૮/૨૦૨૧	૬	૬	૫
	૨૬/૦૮/૨૦૨૧	૮	૭	૭
	૨૭/૦૮/૨૦૨૧	૭	૭	૭
	૨૮/૦૮/૨૦૨૧	૮	૮	૮
	૨૯/૦૮/૨૦૨૧	૮	૮	૭