

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/324169183>

VARIETAL SCREENING OF VARIETIES/ BREEDING MATERIALS FOR RESISTANCE TO DIFFERENT DISEASES IN NATURAL CONDITION UNDER SOUTH GUJARAT REGION

Article in *International Journal of Current Advanced Research* · August 2017

DOI: 10.24327/ijcar.2017.5660.0772

CITATIONS

0

READS

61

1 author:



Rakesh K. Patel

Navsari Agricultural University

15 PUBLICATIONS 3 CITATIONS

SEE PROFILE



VARIETAL SCREENING OF VARIETIES/ BREEDING MATERIALS FOR RESISTANCE TO DIFFERENT DISEASES IN NATURAL CONDITION UNDER SOUTH GUJARAT REGION

Prashant B. Sandipan., Bhanderi G. R., Patel R. K and Solanki B.G

Main Cotton Research Station (MCRS), Navsari Agricultural University (NAU), Surat 395 007 (Gujarat), India

ARTICLE INFO

Article History:

Received 25th May, 2017

Received in revised form 19th

June, 2017 Accepted 14th July, 2017

Published online 28th August, 2017

Key words:

Cotton, Screening, Bacterial leaf blight, Alternaria, Diseases, Resistance

ABSTRACT

Cotton is the main principal commercial crop of our economy. Cotton leftovers the most phenomenal fiber crop and is still nature's wonder fiber. Cotton is grown all around the globe, and is traded internationally as well. The production is inclined by the repeated out breaks of pest and diseases and these are the major factors responsible for lower yield of cotton. A field trial was conducted at Main Cotton Research Station (MCRS), NAU, Surat in the year 2014 for a season to determine the influence of diverse diseases in respect to different varieties/ entries. During the season, entries/ varieties were evaluated for their reaction against the different diseases in cotton crop. Total thirty six varieties/ entries including checks were evaluated for their reaction against the bacterial blight disease. Out of these entries, GSHV 159 and GSHV 172 found disease free. Whereas, eleven entries viz., GSHH 2529, GSHH 2646, GTHH 193, G. Cot. Hy.8, GISV 272, GBHV 164, GBHV 180, G. Cot. 16, G. Cot. 20, G. Cot. 22 and GBHV 185 were found moderately resistant and G. Cot. Hy. 12 and G. Cot. 10 were moderately susceptible and rest of the entries was found resistant against the bacterial leaf blight disease.

Copyright©2017 Prashant B. Sandipan et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

INTRODUCTION

Cotton is a main commercial of our country and natural fibre know as "White Gold" contains with soft, fluffy staple fiber that grows in a boll, or protective case, around the seeds of the cotton plant. The fiber is almost pure cellulose. The plant is a shrub native to tropical and subtropical regions around the world, including the Americas, Africa and India.

It is a white fibrous agricultural product that has a wide variety of uses, from textile production, to creating paper, to producing oil and food products. Cotton is grown all around the globe, and is traded internationally as well. Higher production of cotton requires high investment in terms of fertilizers, pesticides, weedicides and other cultural operations. This ultimately sometimes leads to or influenced by the repeated out breaks of pest and diseases and occasionally responsible for lower yield of cotton. Out of 30 diseases known to occur in cotton crop from time to time, the bacterial blight is the most wide spread and destructive disease reported to cause yield losses of about 10 to 30 per cent (Bhatti and Bhutta, 1983, Kalpana et al., 2004, Sekhon et al., 2008 and Sandipan et al., 2016) and also affect the quality of lint (Sharma and Chauhan, 1985). Bacterial leaf blight, boll rots, wilts and leaf spots are the most destructive cotton diseases (Chopra, 1977). Under natural, bacterial blight

infection, boll yield losses up to 35 % have been reported (Sheo Raj and Verma, 1988). Leaf spots rank third among the diseases on cotton in India. Among the leaf spots, bacterial blight (*Xanthomonas campestris* pv. *malvacearum* (Smith), Alternaria leaf spot (*Alternaria macrospora* Zimm) and grey mildew (*Ramularia aerea*) have been reported to be damaging. Bacterial leaf blight (BLB) of cotton caused by *Xanthomonas campestris* pv. *malvacearum* (Smith) Dye affects the entire aerial parts of cotton plant i.e. necrosis of parenchymatous tissue in the local phase and blockage of xylem vessels in its systemic phase (Casson et al., 1977). In north India, the cotton leaf curl virus disease (CLCuD) caused by a Gemini virus and transmitted by whitefly, *Bemisia tabaci* has become a major threat to cotton cultivation since its appearance in 1993 (Monga et al., 2011). Seed is the costliest input and is highly prone to losses in germination and vigour due to seed mycoflora. Seeds acts as carrier in transmission of pathogens and thereby causes economic threat to cotton cultivation. Resistant varieties are the valid option and a good source in any disease management strategies. Considering this whole scenario, this present exploration was made to ascertain the spectrum of fungal flora associated with the cotton seeds.

Word cloud from the Introduction

Agro meteorological situation of MCRS, Surat (Gujarat)

In this year, monsoon was delayed in Surat about one month, after that there was sufficient rainfall during the month of July-14. Sowing of most of the experiments was completed

*Corresponding author: Prashant B. Sandipan

Main Cotton Research Station (MCRS), Navsari Agricultural University (NAU), Surat 395 007 (Gujarat), India

manage the diseases in the long run. Screening was therefore undertaken to evaluate a number of cotton entries against the bacterial blight disease and other diseases during *kharif* 2014. Total thirty six varieties/ entries including checks were evaluated for their reaction against the bacterial blight disease. Out of these entries, GSHV 159 and GSHV 172 found disease free. Whereas, eleven entries *viz.*, GSHH 2529, GSHH 2646, GTHH 193, G. Cot. Hy.8, GISV 272, GBHV 164, GBHV 180, G. Cot. 16, G. Cot. 20, G. Cot. 22 and GBHV 185 were found Moderately resistant and G. Cot. Hy. 12 and G. Cot. 10 were Moderately susceptible and rest of the entries were found resistant against the bacterial blight disease. Moreover, other diseases were not observed.

Reaction of cotton varieties/cultures against bacterial blight disease at MCRS, NAU, Surat during 2014-15

Sr. No.	Entries	Bacterial leaf blight		
		PDI	Max Grade	Reaction
MLT hirsutum - Hybrids				
1	GSHH 2529	2.0	2	MR
2	GSHH 2646	1.5	2	MR
3	GSHH 2777	0.5	1	R
4	GTHH 193	2.5	2	MR
5	GTHH 194	0.5	1	R
6	GTHH 209	0.5	1	R
7	GTHH 49	0.5	1	R
8	G. Cot. Hy.8	10.0	2	MR
9	G. Cot. Hy.12	11.5	3	MS
MLT of <i>G. hirsutum</i>				
10	GSHV 159	0.0	0	DF
11	GISV 216	0.5	1	R
12	GISV 272	5.0	2	MR
13	GBHV 164	6.0	2	MR
14	GBHV 170	1.0	1	R
15	GBHV 177	2.5	1	R
16	GBHV 180	6.5	2	MR
17	G. Cot. 10	12.5	3	MS
18	G. Cot. 16	9.0	2	MR
19	G. Cot. 20	8.5	2	MR
20	G. Cot. 22	10.5	2	MR
LSVT of <i>G. hirsutum</i>				
21	G. Cot. 10	14.5	3	MS
22	GSHV 172	0.0	0	DF
23	GSHV 173	1.5	1	R
24	GSHV 175	4.5	1	R
25	GBHV 12	3.0	1	R
26	GBHV 183	3.5	1	R
27	GBHV 185	7.5	2	MR
28	GJHV 473	0.5	1	R
29	GJHV 517	2.0	1	R
30	GJHV 519	4.0	1	R
31	GJHV 526	6.5	1	R
32	GTHV 8/10	4.5	1	R
33	GTHV 10/25	3.0	1	R
34	GTHV 10/28	3.5	1	R
35	G. Cot. 16	9.5	2	MR
36	G. Cot. 20	12.0	2	MR

Acknowledgement

Author is highly thankful to Main Cotton Research Station (MCRS), NAU, Surat (Gujarat) for providing the required facility and other arrangements for the conducting the experiment.

References

Anil, G. H. 2013. Studies on leaf blight of Bt cotton caused by *Alternaria* spp. M.Sc. thesis submitted to the University of Agricultural Sciences, Dharwad.

Bhatti, M. A. R. and Bhutta, A. R. 1983. Incidence of seed borne bacterial infection in different cultivars of cotton. *Pakistan Cottons*, 27: 73-74.

Casson, E. T., Richardson, P. E., Brinkerhoff, L. A. and Gholson, R. K. 1977. Histopathology of immune and susceptible cotton cultivars inoculated with *Xanthomonas campestris* pv. *malvacearum*. *Phytopathology*, 67: 195-196.

Chopra, B. L. 1977. Important diseases of cotton and their control measures in India. Paper presented at staff course on cotton production, its processing and marketing technology held at P.A.U., Ludhiana.

Hosagoudar, G. N., Chattannavar, S. N. and Kulkarni, S. 2008a. Screening of Bt and Non Bt cotton genotypes for foliar diseases. *Karnataka J. Agric. Sci.*, 21 (1): 141-143.

Hosagoudar, G. N., Chattannavar, S. N. and Kulkarni, S. 2008b. Survey for foliar diseases of Bt cotton. *Karnataka J. Agric. Sci.*, 21 (1): 139-140.

Hussain, T. and Tahir, M. 1993. Chemical control of bacterial blight of cotton. *Pak. J. Phytopathol.* 5 (1-2): 119-121.

Kalpana, P., Chellamuthu, V. and Jeyalakshmi, C. 2004. Screening of cotton hybrids against bacterial blight incited by *Xanthomonas campestris* pv. *malvacearum* (Smith) Dye, Paper presented in Inter. Symp. Strat. Sust. Cotton Prod. - A Global Vision 3, Crop Production, 23-25 November 2004, Univ. Agric. Sci., Dharwad (India), pp. 373-374.

Khan, M. A. and Ilyas, M. B. 1999. Cotton germplasm response of slow blighting against *Xanthomonas campestris* pv. *malvacearum* and slow curling against CLCuV infection. Proc. 2nd. National Conf. of Pl. Path. Sep. 27-29, U.A.F. pp. 138-139.

Lopez-Lavalle, LAB, Potter N., Brubaker, C. L. 2012. Development of a rapid, accurate glasshouse bioassay for assessing fusarium wilt disease responses in cultivated *Gossypium* species. *Plant Pathol.* 61: 1112-1120.

Monga, D., Kranthi, K. R., Gopalakrishnan, N., Mayee, C. D. 2011. Changing scenario of cotton diseases in India - the challenge ahead. , 2011 pp. 272-280.

Sandipan B Prashant, Desai H. R. and Solanki B. G. (2016). Cotton Pathology. In: Cotton Research in Gujarat, Technical Bulletin, University publication No.88/2015-16, P. 100 published by Research Scientist (Cotton), MCRS, NAU, Surat (Gujarat).

Sharma, B. K. and Chauhan, M. S. 1985. Studies on the chemical control of foliar diseases of cotton in Haryana state. *Agric. Sci. Digest.* 5: 153-56.

Sheo Raj and Verma, J. P. 1988. Diseases of cotton in India and their management. Review of Tropical Plant Pathology, 5: 207-254.

Sekhon, P. S., Singh, D. and Singh, G. 2008. Cotton pathology research achievements and targets in Punjab. All India Coordinated Cotton Improvement Project Report. PAU, Ludhiana.

Ulloa, M., Huttmacher, R.B., Davis, R.M., Wright, S.D., Percy, R., Marsh, B. 2006. Breeding for Fusarium wilt race 4 resistance in cotton under field and greenhouse conditions. *J Cotton Sci* 10: 114-127.

Wang, C. and Roberts, P. A. 2006. A Fusarium wilt resistance gene in *Gossypium barbadense* and its effect

Varietal Screening of Varieties/ Breeding Materials For Resistance to Different Diseases in Natural Condition Under South Gujarat Region

- on root-knot nematode- wilt disease complex. *Phytopathology* 96: 727-734.
- Wang, P. Z., Su, L., Qin, L., Hu, B. M., Guo, W. Z. and Zhang, T. Z. 2009. Identification and molecular mapping of a Fusarium wilt resistant gene in upland cotton. *Theor Appl Genet.* 119:733-739.
- Wiles, A. B. 1963. Comparative reactions of certain cottons to Fusarium and Verticillium wilts. *Phytopathology* 53:586-588.
- Wheeler, B. E. J. 1969. An Introduction to Plant Disease. John Willey and Sons, London, pages: 374.
- Wu, Z. B., Li, D. B. and Chen, P. 2003. Identification of cotton resistance to Fusarium and Verticillium wilts and the influence of the diseases on yield. *Acta Agron Sin* 29: 947-950.

How to cite this article:

Prashant B. Sandipan *et al* (2017) ' Varietal Screening of Varieties/ Breeding Materials For Resistance to Different Diseases in Natural Condition Under South Gujarat Region ', *International Journal of Current Advanced Research*, 06(08), pp. 5657-5660. DOI: <http://dx.doi.org/10.24327/ijcar.2017.5660.0772>
