

: Research Recommendations:

A) Crop improvement: Varietal Development

1. Little Gourd: GNLG-1 (2012)

It recorded 15.6 tones/ha fruit yield with yield advantage of 32.9 % over local check. It produced more number of fruits per vine in addition to its better quality. In disease and pest reaction, it is also found superior with respect to disease viz., anthracnose, powdery mildew and vine borer as compared to local check.



2. Pointed Gourd: GNPG-1 (2014)

This variety was selection from local germplasm. It has recorded 47.13 % higher fruit yield (15.11 t/ha) over the local variety. The variety has long, light green fruit with fair whitish strip.



3. Brinjal: GNRB 1 (Gujarat Navsari Round Brinjal 1) [2016]

GNRB 1 (Gujarat Navsari Round Brinjal 1) Variety registered 23 % fruit yield superiority over GJB-3 and GOB-1. Calyx is medium size, spineless and purple in colour. Fruits are round, dark purple in colour and have purple green leaves. GNRB-1 had low incidence of little leaf disease reaction (3.90 %) and shoot borer (3.35 %) GNRB-1 is recommended for general cultivation in brinjal growing areas of South Gujarat.



Varieties Endorsed: Sweet Potato: C-71, Cross-4, Bhu Kanti



C-71 (1994-95)



Bhu Kanti (2017-18)

4. Tomato: GT 7

Tomato genotype NTL-12-01 (301.0 q/ha) performed well under South Middle and North Gujarat regions where, it exhibited overall 28.47, 26.54 and 25.82 % increased fruit yield over standard checks viz; JT-3 (234.3 q/ha), AT-3 (226.8 q/ha) and DVRT-2 (228.1 q/ha), respectively. The genotype showed less damage by fruit borer, whitefly as well as leaf miner as compared to standard checks. The genotype NTL 12-01 is recommended for cultivation of farmers of South, North and Middle Gujarat regions as GT 7.



Greater yam: GGY 1: Hemlata (Year of release: 2019)

- This variety had recorded 18.48 t/ha average tuber yield. This is a bio-fortified variety contains purple flesh tuber and rich in total soluble sugar, crude fibre, anthocyanin, phosphorus, potassium, ferrous, zinc and copper content and low in anti-nutritional factor Diosgenin. It showed moderately resistant to anthracnose disease. Recommended for cultivation in Gujarat as “GGY1 (Gujarat Greater Yam-1): Hemlata”.



Release proposal for okra variety NOL-17-05 [Purna Rakshak]

Okra growing farmers of south Gujarat region are advised for cultivation of okra variety NOL-17-05 (GNO-1: Purna Rakshak). The average fruit yield of okra variety NOL-17-05 (GNO-1: Purna Rakshak) is 12.72 t/ha. It exhibited overall yield advantages of 10.70 %, 13.52 % and 12.59 % in *Kharif* season over the check varieties GAO-5, Pusa Sawani and GO-6, respectively. It matures within 89-112 days (medium group), having good in fruit size and plant structure. It has high yield potential and moderately resistant against YVMV, Powdery mildew, ELCV disease as well as moderately resistant against pod borer, jassid and whitefly.



Elephant foot yam: GEFY-1 (Swagata) (Year of release: 2020)

- This variety had recorded 44.84 t/ha mean corm yield. Its light orange fleshed corm is reported to have appreciable amount of starch, dietary fiber, carbohydrate content, protein, vitamin A, iron, manganese, zinc and calcium. The acidity feels same like “Gajendra” while consumption. Resistant reaction against collar rot disease. Recommended for elephant foot yam growing areas of Gujarat as GEFY-1 (Swagata).



Gujarat Tannia 1 (G. Tannia 1: Navsari Pari)

The farmers of Gujarat state are recommended to grow Gujarat Tannia 1 (G. Tannia 1: Navsari Pari) during *kharif* season. The proposed variety recorded average green leaves yield of 7.96 t/ha and after 270 days of planting, corm and cormel yield 10.02 t/ha in south Gujarat which was 31.1 and 14.8 per cent higher over national check variety Konkan Haritparni, respectively. The variety also having higher starch (8.15%) and low fibre content (1.65%) with value added traits. This variety has found lower population of aphid as well as less prevalence of *phytophthora* leaf blight and corm rots as compared to check.



B) Production Technology:

Year: 1985

1. Standardization of fertilizer dose and spacing in tapioca:

The tapioca variety H165 should be planted in first week of April at 90 cm x 90 cm spacing. The crop should be manured at the rate of 12.5 tons FYM/ha as basal while land preparation and fertilized at the rate of 75:75:75 kg/ha N: P₂O₅K₂O for higher production of tubers. The fertilizers are to be applied in two equal splits viz. first half at the time of planting and the remaining half at two months after planting.

Year: 1990-91

1. Standardization of fertilizer dose in cabbage:

The maximum production in cabbage var. Pride of India could be obtained through application of 75 kg N/ha in two equal splits. The first half of N should be applied at the time of transplanting and the second half at 30 days after transplanting.

2. Standardization of seed production technology in cauliflower cv. Early Kunwari:

Different methods for production of cauliflower seeds var. Early Kunwari were tried and the seed to seed method involving raising nursery during last week of August and transplanting during last week of September was found best. Further, maximum seed production (517 kg/ha) could be obtained by applying 120 kg N/ha in two equal splits viz. first at transplanting and second one month after transplanting.

Year: 1992-93

1. Sprinkler irrigation in cabbage:

The crop of cabbage could be irrigated through sprinkler at 11 to 14 days interval keeping 5 cm depth. The sprinklers should be operated at 2.75 kg/cm² pressure with an application rate of about 1.67 cm/hr for about 3 hours.

2. Sprinkler irrigation in cow pea:

The crop of cowpea cv. PusaPhalguni could be irrigated through sprinkler at 9 to 10 days interval up to March and 7 to 8 days interval during April and May keeping 5 cm depth. The sprinklers should be operated at 2.75 kg/cm² pressure with an application rate of about 1.67 cm/hr for about 3 hours.

Year: 1993-94

1. Standardization of fertilizer dose in cauliflower cv. Pusa Deepali :

The winter crop of cauliflower cv. Pusa Deepali should be fertilized at the rate of 40:40 kg N:P₂O₅ per hectare as basal dose followed by an application of 40 kg N/ha at 30 days after T.P.

2. Sprinkler irrigation in cauliflower:

The crop of cauliflower could be irrigated through sprinkler at 11 to 14 days interval keeping 5 cm depth. The sprinklers should be operated at 2.5 kg/cm² pressure for about three hours.

3. Standardization of sowing time, spacing and fertilizer dose in Indian Bean for summer planting

Indian bean cv. Kapasi should be sown in the third week of May at distance of 60 cm between the rows and 30 cm within the row for higher production of green pods. The

crop need 60 kg N/ha to be applied in two equal splits; viz. first half as basal and the remaining half at 20 days after sowing.

Year: 1995-96

1. Standardization of fertilizer doses for okra cultivation:

The Okra crop cv. Parbhani Kranti should be fertilized at rate of 150 kg N/ha in two equal splits. The first half dose of nitrogen (75 kg/ha) and full dose of P and K each @ 50 kg/ha should be applied as a basal and the remaining 75 kg N/ha should be applied at 45 days after sowing.

2. Sprinkler irrigation in okra:

The crop of okra could be irrigated through sprinkler. Under enough water availability, the sprinkler system should be operated for three hours at ten days interval. Under limited water availability conditions, the system should be operated for three hours at 18 days interval. The sprinklers should be operated at 2.75 kg/cm².

Year: 1997

1. Standardization of fertilizer doses for okra cultivation:

The okra crop cv. Parbhani Kranti should be fertilized at the rate of 20 t/ha with seasoned press mud 15 days prior to sowing and 75 kg N/ha in three equal splits each at sowing and 30 and 60 days after sowing.

2. Standardization of fertilizer doses for onion cultivation:

The maximum production in onion cv. Pusa Red could be obtained through application of 125 kg N/ha in two equal splits. The first half dose of nitrogen (62.5 kg/ha) and full dose of P and K each @ 50 kg/ha should be applied as a basal and the remaining half dose of nitrogen should be applied at 30 days after transplanting.

Year: 1998

1. Standardization of fertilizer doses for cabbage cultivation:

The cabbage crop cv. Golden Acre should be fertilized at the rate of 200 kgN/ha in two equal splits. The first half dose of nitrogen (100 kg N/ha) and full dose of P and K @ 50 kg/ha should be applied as a basal and the remaining 100 kg N/ha should be applied at 30 days after transplanting.

2. Standardization of fertilizer doses and spacing for capsicum cultivation:

The crop of capsicum cv. California Wonder should be transplanted at a distance of 45 x 20 cm spacing. The crop should be fertilized at the rate of 120:50:50 kg/ha NPK for higher production. The first half dose of nitrogen (60 kg/ha) should be applied as basal along with P and K each at 50 kg/ha while transplanting and the remaining half dose of nitrogen to be applied at 30 days after transplanting.

Year: 1999

1. PGRs application in capsicum:

The maximum production in capsicum Cv. California Wonder could be obtained through one spray of NAA-10 ppm at flower initiation stage.

Year: 2000**1. PGRs application in okra:**

The farmers are advised to spray summer Okra Var. Parbhani Kranti with NAA 75 mg/l at 15 and 30 days after sowing to get higher yield of fruit and seed.

2. Standardization of fertilizer doses and spacing for bottle gourd cultivation:

The farmers are advised to sow bottle guard cv. Pusa Naveen at 2.0 X 1.0 m distance and should be fertilized with 50 kg N, 50 kg P₂O₅ and 50 kg K₂O along with 10 ton FYM per hectare as basal dose followed by 50 kg N/hectare at 30 days after sowing.

Year: 2002**1. Standardization of fertilizer doses and spacing in water melon:**

The farmers of South Gujarat are advised to sow Watermelon cv. Sugar Baby in single row at 2x1 m distance or in paired row at 3.4 x 1 x 0.6 m distance and fertilize it with 150 kg N, 50 kg P and 50 kg K along with 20 t FYM per hectare.

Year: 2007**1. Standardization of fertilizer doses for brinjal cultivation:**

It is recommended to the peasantry of South Gujarat area, growing brinjal cv. "SuartiRavaiya" during *rabi* season, to fertilize their brinjal plant with the combination of 75 % recommended dose of fertilizer (75:28:28 kg N, P₂O₅, K₂O) along with 20 tones of Press-mud/ ha or 10 tones of Bio-compost/ ha, to obtain higher yield as well as to improve the soil health.

Year: 2011-12**1. Effect of land configuration, soil conditioner and fertilizer on greater yam:**

The farmers of south Gujarat heavy rainfall zone, AES-III growing greater yam (*Dioscorea alata* L.) cv. Local Round are advised to plant the crop on ridge furrow of 30 cm height at 90x 90 cm distance and fertilize with FYM @ 20 t/ha along with recommended dose of fertilizer @ 80 : 60 : 80 NPK kg/ha. Full dose of FYM, P₂O₅ and half dose of N and K₂O applied at plating then remaining half dose of N and K₂O should be applied in two equal split at 90 and 135 days after planting for getting higher tuber yield and maximum economic return (1: 2.95).

2. Organic production of elephant foot yam:

The farmers who want to grow elephant foot yam organically are advised to apply either Vermicompost @ 5 t/ha + *Azospirillum* @ 5 kg /ha + *Phosphobacteria* 5 kg/ha + ash @ 5 t/ha or FYM @ 10 t/ha + *Azospirillum* @ 5 kg /ha + *Phosphobacteria* 5 kg/ha + ash @ 5 t/ha).

The farmers of South Gujarat intending to grow elephant foot yam (cv. Gajendra) are advised to apply FYM @ 10 t/ha + 80: 60: 100 NPK kg/ha for getting higher net income of Rs. 2.9 lakh and BCR of 2.9.

3. Effect of different organic manures on growth, yield and quality of yam:

The farmers who want to grow greater yam organically are advised to apply 75% N through vermicompost and 25% N through castor cake for getting net income of Rs. 1.9 lakh and BCR of 2.4. The farmers of South Gujarat intending to grow greater yam are

advised to apply 80-60-80 kg N-P₂O₅-K₂O/ha. For getting higher net income of Rs. 2.8 lakh and BCR of 2.8.

Year: 2012-13

1. Effect of different organic manures on growth, yield and quality of organically grown turmeric (*Curcuma longa*):

The farmers of South Gujarat heavy rainfall zone AES III growing turmeric variety Sugandhum are advised to apply bio compost (1.8% N) or vermi compost (1.2% N) + neem cake (5.1% N) in equal proportion to supply N @ 60 kg/ha for achieving higher rhizome yield with superior quality of turmeric as well as net income. Application of manure in this manner also improves the soil health.

Year: 2013-14

1. Effect of plant density and sett size on growth and dry matter partitioning of elephant foot yam:

The farmers of south Gujarat heavy rainfall agro-climatic zone growing elephant foot yam cv. Gajendra are advised to plant elephant foot yam at the distance of 60 cm × 60 cm by using seed corm sett of 250 g weight for obtaining higher BCR. By this way, farmers can obtain higher yield and save the cost of seed corm.



Yield from 250 g seed corm sett



Yield from control sett

2. Effect of banana pseudostemsap and vermiwashspray on organically grown onion:

The farmers of South Gujarat heavy rainfall zone (AES III) those who want to grow onion cv. Pilipati are advised to apply 125 kg N/ha through 2.4 t/ha biocompost, 0.9 t/ha castor cake and 3.2 t/ha vermicompost in equal proportion of nitrogen along with foliar spray of enrich banana pseudo stem @ 2% spray or 2% spray of banana pseudo stem and enrich banana pseudo stem in the 1:2 ratio to get high net return. Organic manures should be apply at the time of transplanting and one month after transplanting whereas, the liquid manures should be apply at 15 days interval starting from 15 days after transplanting i.e. 15, 30 and 45 days after transplanting.

3. Feasibility of organic farming in tomato cv. Junagadh Tomato -3:

The farmers of South Gujarat heavy rainfall zone (AES III) those who want to grow organic tomato cv. Junagadh Tomato at 60 cm x 60 cm spacing are advised to apply 75 kg N/ha through 2.1 t/ha biocompost (50%N) + 0.9 t/ha castor cake (50%) or 4.6 t/ha vermicompost (75% N) + 0.4 t/ha neem cack (25% N) in two splits (at the time of transplanting and one month after transplanting) to get higher yield and net profit. Common dose of *Azotobacter* biofertilizers @ 2 kg/ha with organic manures at the time of transplanting. To prevent the pest and disease infestation, foliar spray of vermiwash @ 0.5% and cow urine @ 1% at monthly interval after transplanting is beneficial.

Year: 2014-15

1. Integrated Nutrient Management in Little gourd:

The farmers of South Gujarat heavy rainfall zone AES III cultivating little gourd cv. GNLG-1 are advised to follow INM to fertilize the crop as per the schedule given below to get higher better quality fruits and net realization.

Basal dose: Apply 10 t/ha well decomposed FYM, 25 kgN/ha through Bio compost on equivalent N basis along with 50 kg/ha each of P and K by chemical fertilizer.

Top dressing: Apply 25 kg N/ha in two splits through chemical fertilizer at 30 and 60 days after Planting

Note: 1. In subsequent years, apply fertilizer as above schedule.

2. Pruning should be done in month of December.

2. Effect of different organics on growth and yield of brinjal cv. SurtiRavaiya(pink):

The farmers of South Gujarat heavy rainfall zone AES III intended to grow brinjal variety SurtiRavaiya (Pink) organically are advised to apply castor cake (4.5 % N ; dry weight basis) in two equal proportion to supply N @ 100 kg/ha for achieving higher yield and net income as well as to improve the soil health.

Apply 4.5 t/ha castor cake in two equal splits at the time of transplanting and one month after transplanting.

Note :

– *Trichoderma virides* should be applied at the rate of 5 kg/ha at the time of transplanting.

– Maize should be grown as trap crop on the border.

– Sticky trap should be used @ 40/ha.

– Tricho card should be used @ 5/ha.

After transplanting apply foliar spray of neem based pesticide and cow urine at monthly intervals.

3. Response of seed sowing on germination, growth, flowering and yield of Spine gourd (*Momordica dioica*Linn.) cv. Local:

The farmers of South Gujarat heavy rainfall zone AES-III interested to grow spine gourd cv. Local through seed are advised to sow five seeds per dibble on raised bed in last week of March and mulch with paddy straw for higher fruit yield.



4. Performance of greater yam (*Dioscorea alata* L.) under different stacking systems:

The farmers of south Gujarat heavy rainfall zone AES III growing greater yam cv. Local Round are advised to plant greater yam at the distance of 90 cm × 90 cm with elephant foot yam cv. Local as a live stacking crop in-between two rows of greater yam at a distance of 90 cm × 90 cm and train the vines of greater yam on the plants of elephant foot yam with application of 15 tonne of FYM and 120:90:120 kg NPK/ha to obtain higher yield and net return.



5. Effect of rates of castor cake and Banana Pseudostem sap on yield and quality of organically grown Garlic (*Allium sativum* L.):

The farmers of South Gujarat heavy rainfall zone AES III growing garlic organically are advised to apply recommended 100 kg N/ha through organic manures as per schedule given below to get higher yield and net profit.

- Apply 1.4 t/ha biocompost and 3.3 t/ha vermicompost at the time of sowing and 0.7 t/ha castor cake one month after sowing.
- Apply 2000 lit/ha banana pseudostem sap at 35 and 55 days after sowing

Note:

- Apply common dose of *Azotobacter* biofertilizer @ 2 kg/ha.
- After sowing, apply foliar spray of neem based insecticide and cow urine at monthly interval.
- Maize should be grown as trap crop at the border.
- Sticky trap should be used @ 40/ha.

Year: 2015-16

1. Standardization of fertigation and methods of training in cucumber under naturally ventilated poly house:

Train plants to single stem system. Fertigate the crop with 9.0:7.5:7.5 kg NPK along with application of 0.5 kg *Trichoderma viride*, 0.5 litre *Pseudomonas fluorescens*, 2 t FYM or 0.4 t vermicompost and 5.0 kg micro-nutrient (Grade-5) at the time of sowing for higher net returns.



2. Site specific nutrient management study of Elephant foot Yam

The soil having deficient N and sufficient P and K then apply 100: 45: 75 NPK kg/ha in two splits. First dose of 50: 45: 37.5 NPK kg/ha at 45 days after planting. Second dose of 50: 00: 37.5 NPK kg/ha one month after application of first dose.



Year: 2016-17

1. Effect of rhizome size on growth and yield of turmeric cv. GNT-1.

The farmers of south Gujarat heavy rainfall zone are advised to plant mother rhizome pieces (10-15 g) of turmeric cv. GNT-1 in pro tray and transplant it after one month in field with minimum quantity of seed rhizomes.



2. Standardization of fertigation and methods of training in capsicum under naturally ventilated polyhouse.

Farmers cultivating capsicum in naturally ventilated polyhouse (1000 m² area) are advised to fertigate the crop with 25: 25: 25 kg NPK along with application of 0.5 kg *Trichoderma viride*, Phosphorous Solubilizing Bacteria (*Bacillus megaterium*), Azotobactor, *Pseudomonas fluorescens* each, 0.4 t vermicompost and 5.0 kg micro-nutrients (Grade-5) at the time of planting and train plants to four shoot system for higher net returns.



Year: 2017-18

1. Integrated Nutrient Management in cauliflower (*Brassica oleracea* var. botrytis):

The farmers of South Gujarat Agro climatic Zone-I growing cauliflower are advised to apply 20 kg N + 40 kg P₂O₅ along with 20 t/ha FYM and 5.70 t/ha Bio compost as basal doze. The 20 kg Nitrogen should be applied 30 DAT as top dressing to get higher yield and return.

2. Response of okra to foliar application of Silicon

The farmers of South Gujarat growing summer okra are advised to spray silicon based liquid fertilizer @ 2ml /l (silicon: 0.79% w/v + boron :0.18% w/v- OSAB – Si⁺) at 30,45 and 60 DAS to obtain higher yield and net income.



Effect of Foliar application of silicon on okra.

3. Performance of grafted vs. non-grafted brinjal during rainy season under South Gujarat conditions

The farmers of South Gujarat Heavy Rainfall Zone-I (AES-III) are advice to adopt grafting technique using wild species (*Solanum torvum*) as rootstock and pink and purple *Surati Ravaiya* brinjal as scion for better plant survival during rainy season, better fruit set, comparatively less shoot and fruit borer infestation, extended life span, higher yield and net returns.





4. Comparative performance of different parthenocarpic cultivars of cucumber through vegetative propagation under polyhouse conditions.

Farmers cultivating parthenocarpic cucumber varieties in greenhouse are advised to use newly pruned side shoots of current crop as propagating material for raising of successive crop without paying high price for seed which performs equally well to the crop raised from seeds and concurrently, excessive plants generated from pruned side shoots can be sold for additional income.





Year: 2019-20

1. Effect of different sources of nutrients and fertigation levels on yield and other horticultural traits in tomato under protected culture

Farmers cultivating tomato in naturally ventilated polyhouse (1000 m²) are advised to fertigate the crop with 25: 12.50: 12.50 kg NPK (As per the schedule given in table below) through water soluble fertilizers along with application of 0.5 kg *Trichoderma viride* and *Pseudomonas fluorescens* each, 0.5 L Phosphorous Solubilizing Bacteria (*Bacillus megaterium*) & potash mobilizer- *Frateuria aurantia* each, 2 t FYM and 5.0 kg micro-nutrients (Grade V) at the time of transplanting for higher yield as well as net returns.



2. Feasibility of tomato cultivation through grafting during rainy season

The farmers of South Gujarat Heavy Rainfall Zone-I are advised to adopt interspecific grafting of tomato with *Solanum torvum* for better plant survival during rainy season, extended life span, more number of fruits, comparatively less leaf curl

infection, white fly, leaf miner, fruit borer infestation, higher yield and net returns



3. Artificial oscillation for increasing fruit set and performance of tomato in polyhouse under South Gujarat conditions

Farmers cultivating tomato in naturally ventilated polyhouse are advised to vibrate tomato truss with electric pollinator on every 3rd day starting from the day of first flowering for 10 seconds during morning hours between 7.30 am to 9.00 am for better fruit set, higher yield and net returns.



4. Integrated Nutrient Management in cabbage (*Brassica oleracea* var. *capitata*)

The cabbage growing farmers of south Gujarat are advised to grow cabbage under INM system and fertilize their crop with combination of 50% recommended dose of nitrogen (200:00:37.5 NPK kg/ha) along with Bio compost (Nitrogen based) to obtain higher yield and income. Entire quantity of Bio compost and potash as well as half quantity of nitrogen should be applied as basal. Remaining half dose of nitrogen should be applied as top dressing in two equal splits viz., 30 and 45 DATP.

5. Validation of organic farming technologies in elephant foot yam

The farmers of South Gujarat Heavy Rainfall Zone, intending to grow elephant foot yam cv. Gajendra organically are recommended to use organic treatment as per below mentioned management:

- Raise green manure of cowpea with 20 kg ha⁻¹ seed rate and incorporate it at 45-60 days before planting of elephant foot yam.
- Take organically produced planting material of 500 g weight and treat it with bucket full of cow dung slurry containing 1- 2 kg neem cake and *Trichoderma harzianum* (5 g per kg seed corm) and then dry under shade before planting.
- At the time of planting, apply FYM : neem cake mixture (in 10:1 ratio) @ 36 t ha⁻¹ incorporated with *Trichoderma harzianum* @ 2.5 kg per tonne of FYM neem cake mixture along

with neem cake @ 1 t ha⁻¹ in pits.

- Raise green manure cowpea again with 20 kg ha⁻¹ seed rate in-between fallow space of elephant foot yam plants and incorporate at 45-60 days in pits along with 3 t ha⁻¹ of ash.



Year: 2020-21

1. Artificial oscillation for increasing fruit set and performance of tomato in polyhouse under South Gujarat conditions

Farmers cultivating tomato in naturally ventilated polyhouse are recommended to vibrate tomato truss with electric pollinator on every 3rd day starting from the day of first flowering for 10 seconds during morning hours between 7.30 am to 9.00 am for better fruit set, higher yield and net returns.



2. Standardize the fertilizer dose of drumstick (*Moringa spp.*)

The farmers of south Gujarat agro climatic zone growing drumstick are recommended to apply 100-75-50 g NKP (217.3 g NCU - 468.7 g SSP - 83.3 g MOP) per tree to obtain higher yield and income. Nitrogen (Neem Coated Urea) apply in four equal splits, first apply at pit preparation (common application of 8 kg FYM enrich with (*Azotobactor*+ *PSB*+ *KMB* (each @ 2ml) in a pit), along with half dose phosphorus (Single Super Phosphate) and potash (Muriate of Potash), second fertilizer application after 30 days after planting; third split application was given after six month interval after pruning along with half dose of phosphorus and potash and fourth split application was given 30 days after pruning.

Application time	Fertilizer	Quantity
First	25 g N ₂ 37.5 g P ₂ O ₅ 25 g K ₂ O	54.30 g NCU+ 234.35 g SSP + 41.65 g MOP
Second	25 g N ₂	54.30 g NCU
Third	25 g N ₂ 37.5 g P ₂ O ₅ 25 g K ₂ O	54.30 g NCU+ 234.35 g SSP + 41.65 g MOP
Forth	25 g N ₂	54.30 g NCU

Note: Common application of 8 kg FYM enrich with (*Azotobactor*+ *PSB*+ *KMB* (each @ 2ml) in a pit



3. Effect of IBA and number of nodes on stem cutting on propagation of little gourd

The farmers/nurserymen of south Gujarat agro climatic zone are recommended to multiplication of little gourd are recommendation to select one year old little gourd vine cutting with two nodes dipped in 80mg/l IBA solution for 30 minute and put in growing media Soil: FYM: Sand (1:1:1) increases sprouting percentage, vine length, vine diameter, root length, root diameters and survival percentage of little gourd cutting.



4. Response of okra to foliar application of Novel Organic Liquid Nutrients and Micronutrients

The okra growing farmers of South Gujarat Agro-Climatic Zone are recommended for foliar application of 1.5 % Novel Organic Liquid Nutrients at 30, 45 & 60 DAS to obtained higher yield of okra.



Year: 2021-22

1. Response of okra to foliar application of Novel Organic Liquid Nutrients and Micronutrients

Farmers of south Gujarat growing *kharif* okra are recommended to apply foliar spray of 1.5 % (150ml/10 litre water) Novel Organic Liquid Nutrients at 30, 45 and 60 DAS along with recommended dose of fertilizer (100-50-50 N-P-K kg/ha) to obtain higher yield and net return.



2. Effect of sowing dates and spacing on off season okra

Farmers of south Gujarat are recommended to cultivate off season okra by sowing in 2nd week of October with spacing of 45 cm x 10 cm to obtain higher profit.



3. Effects of boron and molybdenum on nodulation, growth and yield of cowpea (*Vigna unguiculata* L. Walp.).

Farmers of south Gujarat growing summer cowpea are recommended to give seed treatment of molybdenum @ 2mg/l (Ammonium molybdate @2.40 mg/l) for 24 hours prior to sowing followed by foliar spray of boron @ 4mg/l (Boric acid @ 22.88 mg/l) at 30, 45 and 60 DAS to obtain higher pod yield. Moreover, seed treatment of molybdenum 4 mg/l increases nodulation.



Year: 2022-23

1. Effect of organic spray on growth, yield and quality of tomato (*Solanum lycopersicum* L.) under South Gujarat condition.”

Farmers of South Gujarat Agro-climatic Zone are recommended to spray of Novel Organic Liquid Nutrients 1% (100 ml/10 liter of water) at 25, 50, 75 & 100 DATP along with recommended dose of fertilizer (100-50-50 N-P-K kg/ha) for higher yield and net realization in tomato cv.GT-7.



2. Influence of sett size and spacing on growth and yield of greater yam (*Dioscorea alata* L.)

The farmers of South Gujarat agro climatic zone growing greater yam (Hemlata) are recommended to use 250 g tuber size and planting the greater yam at 90 cm x 60 cm spacing to get higher yield and net realization.



For Scientific Community

Year: 2016-17

1. Evaluation of parthenocarpic cultivars of cucumber under protected conditions for yield and other horticultural traits:

Greenhouse cucumber cultivars Oscar and Valleystar were identified as the highest yielders under naturally ventilated polyhouse, which were at par in performance with cvs. RS 03602833, Kian and Multistar. Evaluation of cucumber cultivars for various sensory parameters by heterogeneous panel of evaluators revealed highest overall score in cv. Multistar statistically at par with KUK-9 and 52-23.

2. Evaluation of tomato cultivars under NVPH for yield and other horticultural traits.

Cultivar Bargad was identified as significantly highest yielder with maximum net realization in naturally ventilated polyhouse. Higher number of fruits per plant and minimum occurrence of blossom end rot were observed as major contributing traits towards yield.

Year: 2019-20

1. Effect of different light sources on growth and quality of microgreens.

- Based on the performance of different microgreens for growth parameters like days to first harvest, leaf area (cm^2), fresh weight and quality parameters viz., ascorbic acid, β -carotene, N, P, K, Ca content, total antioxidant activity and overall acceptability under different light sources, electroluminescent light is recommended for growing microgreens inside growing chamber/chamber/room.
- Fenugreek microgreens displayed significantly maximum ascorbic acid, N, Ca content; while beet root and red cabbage revealed maximum β -carotene, K content and antioxidant activity. Based on sensory evaluation, highest score for overall acceptability was obtained by Amaranth microgreens, which was followed by beet root and red cabbage microgreens.

2. Integrated weed management in elephant foot yam

The scientists are informed that spraying of post emergence herbicide-Glyphosate 41 % S. L. 1 kg a.i./ha at 30, 60 and 90 DAP in-between row space of elephant foot yam cv. Gajendra for effective weed management.



Year: 2022-23

Validation of customized fertilizers in sweet potato

It is inform to scientific community that soil application of “customized fertilizer” (325 kg/ha) two times as basal and 1 month after planting as well as foliar application of “micronol sweet potato”(5 ml/lit) three times on 15, 30 and 45 days after planting should be done for getting higher yield in sweet potato.

Customized Fertilizer (%):

N	P	K	Ca	Mg	Zn	B
11	7	11	6	3	0.4	0.1

Micronol sweet potato (%):

Zn	Cu	B	Fe	Mn
2	0.6	0.2	0.5	0.25

